

21957



0440

0300

0420 9500

0420

IN THE U.S. PATENT AND TRADEMARK OFFICE

Inventor Saul TZIPORI et al
Patent App. 10/041,958
Filed 7 January 2002
For HEMOLYTIC UREMIC SYNDROME
Art Unit Not known
Hon. Commissioner of Patents
Washington, DC 20231


PRELIMINARY AMENDMENT

Prior to examination of the above-identified application,
please amend as follows:

IN THE SPECIFICATION

Please replace page 1 of the specification with revised
page 1 enclosed herewith.

Respectfully submitted,
The Firm of Karl F. Ross P.C.



Jonathan Myers, Reg. No. 26,963
Attorney for Applicant

er
24 January 2002
5676 Riverdale Avenue Box 900
Bronx, NY 10471-0900
Cust. No.: 535
Tel: (718) 884-6600
Fax: (718) 601-1099

Enclosure: Revised Specification Page 1

204240 85814001

HUMAN NEUTRALIZING ANTIBODIES AGAINST HEMOLYTIC UREMIC SYNDROME**SPECIFICATION****CROSS REFERENCE TO RELATED APPLICATION**

This application is a continuation-in-part of copending
5 application 09/302,125 filed 29 April 1999 which is a division of
application 08/749,704 filed 15 November 1996, now abandoned.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was made with government support under
AI41326 and DK58993 awarded by the National Institutes of Health.
10 The government has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to new human monoclonal
antibodies capable of neutralizing Shiga or Shiga-like toxins
which cause hemolytic uremic syndrome in mammals, a process for
15 the preparation of the new human monoclonal antibodies and a
method of treating a mammalian subject to prevent the development
of hemolytic uremic syndrome in a mammalian subject by
administering the monoclonal antibodies to the subject. More
particularly the invention relates to human monoclonal antibodies
20 prepared by administering as an antigen to a transgenic mouse
having human genes an inactivated Shiga-like toxin to induce an
immune response, isolating a splenocyte from the transgenic
mouse, fusing the splenocyte to a mouse myeloma cell to form a
hybridoma, and screening human monoclonal antibodies produced by
25 the hybridoma for the ability to neutralize Shiga or Shiga-like
toxins.